

Use of *Pseudomonas fluorescens* as a bioherbicide for cheatgrass and other invasive winter annual grass control

Daniel R. Tekiela, University of Wyoming Assistant Professor and Extension Specialist of Invasive Plant Ecology

Cheatgrass (*Bromus tectorum*) and other invasive annual grasses cover an estimated 100 million acres of the western United States and cover hundreds of thousands of acres in Wyoming.

These grasses are capable of reducing forage quality, biodiversity, and wildlife habitat. There is great interest in effective, long-term control of these grasses. There has been an increased interest in using bioherbicides to manage and control invasive annual grasses.

Work over the past 15 years has identified unique strains of the ubiquitous soil bacteria *Pseudomonas fluorescens* to control invasive annual grasses. These strains were originally identified from wheat fields in Washington with annual grass injury. From those, specific strains were identified that are reported to only affect certain invasive annual grasses. These strains colonize intercellular space in the roots of the target invasive annual grass and reduce plant growth by inhibiting root cell membrane production. This results in stunting of the target plant. There are currently two strains potentially being utilized as bioherbicides, D7 and ACK55.

D7

Pseudomonas fluorescens D7 is registered as a bioherbicide as of 2014 and is being produced by Verdesian Life Sciences, LLC, Cary, NC. It is labeled for control of “Downy Brome (cheatgrass), Medusahead, Japanese Brome, and Jointed Goatgrass on Wheat,

Barley, Triticale, Oats, and Rangeland” (D7 Label); however, some report it has no effect on Japanese brome. D7 is delivered in a freeze-dried form and can be dissolved into a liquid solution and broadcast sprayed or alternatively used as a seed treatment. Because *Pseudomonas fluorescens* is a cool, moisture-loving bacteria, application should occur when temperatures are below 50F in the fall to ensure ideal growing conditions. Lab studies have shown D7 reduced root growth of annual grasses in petri dishes and when tested in growth chambers (Kennedy, Johnson, & Stubbs, 2001). Additionally, three study sites across Washington showed natural populations of cheatgrass had reduced seed production (16-64 percent), shoot mass (0-54 percent), and plant density (0-35 percent) less than a year after application (Kennedy, 1991); however, there has been limited success in duplicating these results. Recent reports suggest results in the field may not be seen for up to three to five years. No published studies examine the long-term effects of D7 beyond this initial year. There is no published evidence of D7’s effectiveness or lack thereof in field conditions in the Rocky Mountain region. Because this is a living organism, environmental conditions are likely far more important for efficacy than for synthetic herbicides, and regionality may be critically important.

ACK55

Pseudomonas fluorescens ACK55 is undergoing the EPA registration process and is not commercially available. ACK55 is reported to be more selective and to only control downy brome (cheatgrass), medusahead, and jointed goatgrass, but is said to more greatly reduce target plant growth. No peer-reviewed information exists on the effects of ACK55, so its effects are largely unknown, especially in the Rocky Mountain region.



MB906

Although also *Pseudomonas fluorescens*, MB906 is not labeled as a bioherbicide but is labeled and marketed as

a soil inoculant to “enhance biodiversity in soil” (MB906 label). This product cannot be used to control annual grasses. Additionally, there is no published information on the effects of this soil inoculant.

STRAIN	REGISTERED AS A BIOHERBICIDE?	SUMMARY OF PUBLISHED EFFICACY DATA
D7	Yes. Labeled for control of downy brome (cheatgrass), medusahead, Japanese brome, and jointed goatgrass. Registered crops include wheat, barley, triticale, oats, and rangeland.	D7 suppressed downy brome in field studies in 1988 (Kennedy et al. 1991), but attempts to duplicate those results in subsequent field studies were largely unsuccessful (Tranel et al. 1993). Lab studies have shown D7 reduces root growth of downy brome and other annual grasses in petri dishes and in growth chambers (Kennedy et al. 2001); however, in those same laboratory studies D7 also caused root growth suppression in some desirable grasses, including wheat, barley, smooth brome, and tall wheatgrass (Kennedy et al. 2001).
ACK55	No. EPA registration as a bioherbicide is pending.	None available.
MB906	No. This product is not registered for use as a bioherbicide for weed control; it is labeled as a soil inoculant to enhance biodiversity in soil.	None available.

RESOURCES

Kennedy, A. C. (1991). Rhizobacteria suppressive to the weedy downy brome. *Soil Science Society of America Journal*, 55(3), 722–727.

Kennedy, A. C., Johnson, B. N., & Stubbs, T. L. (2001). Host range of a deleterious rhizobacterium for biological control of downy brome. *Weed Science*, 49(6), 792–797. [http://doi.org/10.1614/0043-1745\(2001\)049\[0792:HROADR\]2.0.CO;2](http://doi.org/10.1614/0043-1745(2001)049[0792:HROADR]2.0.CO;2)

Tranel, P.J., D.R. Gealy, and A. C. Kennedy. (1993) Inhibition of downy brome (*Bromus tectorum*) root growth by a phytotoxin from *Pseudomonas fluorescens* strain D7. *Weed Technology*, 7:134-139.

Thank you to Jane Mangold (Montana State University), Richard Lee (Bureau of Land Management), Andrew Kniss (University of Wyoming), and William Stump (University of Wyoming).

Keywords: invasive, weed, herbicide, management

B-1296 • January 2017

Issued in furtherance of extension work, acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture. Glen Whipple, director, University of Wyoming Extension, University of Wyoming, Laramie, Wyoming 82071.

Persons seeking admission, employment, or access to programs of the University of Wyoming shall be considered without regard to race, color, religion, sex, national origin, disability, age, political belief, veteran status, sexual orientation, and marital or familial status. Persons with disabilities who require alternative means for communication or program information (Braille, large print, audiotope, etc.) should contact their local UW Extension office. To file a complaint, write to the UW Employment Practices/Affirmative Action Office, University of Wyoming, Department 3434, 1000 E. University Avenue, Laramie, WY 82071.